



PERGAMON

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Planetary and Space Science 51 (2003) 411–414

Planetary
and
Space Sciencewww.elsevier.com/locate/pss

Observations of 2000 DP₁₀₇ in NAOC: rotation period and reflectance spectrum

B. Yang^{a,*}, J. Zhu^{b,a}, J. Gao^{c,a}, H.T. Zhang^a, X.Z. Zheng^a^aNational Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, People's Republic of China^bBeijing Planetarium, Beijing 100044, People's Republic of China^cDepartment of Astronomy, Beijing Normal University, Beijing 100875, People's Republic of China

Received 18 March 2002; accepted 20 January 2003

Abstract

Photometric observations of a Near-Earth-Asteroid (NEA) 2000 DP₁₀₇ were made on four successive nights during October 2000, 1.4–4.7 with the 0.6-m/0.9-m Schmidt telescope of National Astronomical Observatories, CAS (NAOC). The derived rotation period of 0.1156 day was consistent with that obtained by Pravec et al. (IAU Circular No. 7504, 2000). In addition, the relative reflectance spectrum of the asteroid covering 0.35–0.9 μm was obtained with the NAOC 2.16-m telescope adopting a low-resolution grating (10 Å/pix) on October 2, 2000, which revealed that 2000 DP₁₀₇ is an M-type asteroid.

© 2003 Elsevier Science Ltd. All rights reserved.

Keywords: Techniques; Photometric techniques; Spectroscopic-minor planets; Asteroids

1. Introduction

The NEA 2000 DP₁₀₇ was discovered on February 29, 2000 by LINEAR (Blythe et al., 2000) with orbital elements of $a = 1.37$ AU, $e = 0.38$ and $i = 8.66^\circ$. Radar observations of this object made by Ostro et al. (2000) on September 22 and 23, 2000 using NASA's Goldstone radar in Southern California, revealed that 2000 DP₁₀₇ is a binary system—two separate objects travelling at least a kilometer apart at times. Photometric observation of this object made by Pravec et al. (2000) on nine nights during September 25.2–October 4.2 showed a fast variation with a period of 0.11564 ± 0.00001 day with an amplitude of 0.19 mag, and the eclipse/occultation event in the binary system with a period of 1.76 ± 0.02 day. Their measured broad band color indices revealed that 2000 DP₁₀₇ is spectrally flat, consistent with a C-type minor planet.

In this paper, we present the result of our own photometric and spectral observations of 2000 DP₁₀₇ in early October 2000 with the NAOC 0.6-m/0.9-m Schmidt and 2.16-m telescopes.

2. Photometric observation and analysis

2.1. Observation description and data reduction

The photometric observations of 2000 DP₁₀₇ were made on four nights during 2000 October 1–4 with the NAOC 0.6-m/0.9-m Schmidt telescope using a thick 2048 × 2048 CCD with a 58' × 58' field and scale of 1'' .7 pixel⁻¹. The BATC i-filter with a central wavelength of 6660 Å and a band width of 480 Å was used during the observation (Fan et al., 1996; Zheng et al., 1999). The reduction of images was performed using APPHOT, a routine of the Image Reduction and Analysis Facility (IRAF) package. Magnitudes of the 2000 DP₁₀₇ and comparison stars were extracted from each image using aperture photometry. By convention, we chose 5 comparison stars in each frame. The most stable one with the minimum standard deviation at RA 22^h 11^m 45.57^s, Dec 01° 06'50.7'' with V magnitude 11.226 was adopted as the final standard star to obtain the relative magnitude of the asteroid. The magnitude of the standard star was adopted from the TASS tenxcat catalog (Richmond et al., 2000). The reduced magnitudes of the asteroid were corrected for light travel time and distance (Harris and Lupishko, 1989). The photometric data were analyzed using Phase Dispersion

* Corresponding author.

E-mail address: yangbin@vega.bac.pku.edu.cn (B. Yang).




Fig. 1. Theta statistics of data vs. period.

Minimization (PDM), a routine of the IRAF (Stellingwerf, 1978), to obtain the light curve.

2.2. Result

The observations were made during nights that were mostly not of good quality. Observations on October 1, 2000 yielded a period of 0.1156 day with an amplitude of 0.21 mag (Fig. 1), which was quite consistent with the results of 0.11564 day by Pravec et al. (2000). The deviation of about 0.1 mag among the data is related to the eclipse/occultation event around October 1.66 as mentioned in Pravec et al. (2000). The dashed line in Fig. 2 was fitted by rejecting those points (marked as “x”). Data from other nights suffered from large photometric errors and could not be used to get reasonable results.

3. Spectral observations and analysis

3.1. Observation description and data reduction

Spectral observations were made with the NAOC 2.16-m reflector on 2000 October 25 at the same site, adopting a low-resolution grating centered at 6000 Å with 10 Å/pixel dispersion. Spectral data reduction follows the process of the Small Main-Belt Asteroid Spectroscopic Survey (SMASS) (Xu, 1994; Xu et al., 1995). A solar-analog star 16 Cyg B (Hardorp, 1978) was adopted for obtaining the asteroid’s

relative reflectance spectrum (Luu and Jewitt, 1999). The standard spectrum of 16 Cyg B was adopted from the library of stellar optical spectra by Silva and Cornell (1992). By convention, the reflectance spectrum of 2000 DP₁₀₇ was normalized to 0.55 μm. Some strong absorption bands in the spectra of asteroid and 16 Cyg B were caused by the water and oxygen molecules in the Earth’s atmosphere (Cochran and Barnes, 1981), and this telluric feature was removed as well as possible in the final plot.

3.2. Analysis and result

The reflectance spectrum of 2000 DP₁₀₇ obtained from our observation is almost featureless and slightly reddish over the 0.4–0.89 μm wavelength region (see Fig. 3(a)). The spectral character is in accord with the characteristics of X-type asteroid spectra described by Bus (2002).

Spline3 function was used to fit the spectrum of the asteroid and to obtain 46 points ranging from 0.44 to 0.89 μm with an interval of 0.01 μm. The ‘spectral distances’ between this asteroid with the known spectra of 1341 asteroids from SMASS II (Binzel et al., 2001; Bus, 1999) in the 46-dimensional space were calculated, which is defined as follows:

$$D_x = \sqrt{\sum_{n=1}^k (X_n - Y_n)^2},$$




Fig. 2. Fitted light curve of 2000 DP₁₀₇. Zero phase corresponds to UT 2000 October 1.56.




Fig. 3. (a) Reflectance spectrum of 2000 DP₁₀₇ and (b) spectrum of the asteroid (Binzel et al., 2001) with the closest “spectral distance” to 2000 DP₁₀₇.

where D_x is the ‘spectral distance’ between the unclassified spectrum X and a classified spectrum Y , and n represents the individual channels consisting of the spectrum, where the total number of the channels is k . The calculated results are listed in Table 1, where it is clear that the objects with the closest spectral matches are mostly X-type asteroids including successively No. 1–39 (see Fig. 3(b)). Therefore, we classify the NEA 2000 DP₁₀₇ as a member of “X class” (including the E-, M-, and P-types) (Tholen, 1984,1989).

Recent publications of radar observations of this object provide further information (Margot et al., 2002). They estimate a diameter of 800 m of the primary body and of approximately 300 m of the secondary (Margot et al., 2000). In

Table 1
Asteroids class with the minimum “spectral distances” to the 2000 DP₁₀₇

Classification	Minimum distance	Sequence number	Corresponding asteroid
X-type	0.0705	1	(399) Persephone
C-type	0.1242	40	(551) Ortrud
B-type	0.2487	308	(59) Elpis
S-type	0.2912	398	(1977) Shura
Others	> 0.3300	> 400	

addition, Pravec et al. (2000) give the absolute magnitude $H = 18.08 \pm 0.11$ of this object. Adopting the values

Table 2
Albedos of asteroids with the closest “spectral matches”

Sequence number	Asteroid	P_v	Source	Classification
1	(399) Persephone	0.18	IRAS	M
2	(272) Antonia	0.14	IRAS	M
3	(129) Antigone	0.17	TRIAD	M

mentioned above, we derive a mean V-band albedo of this object of 0.15 by the formula (Bowell et al., 1989)

$$\log P_v = 6.259 - 2 \log d - 0.4H_v,$$

where P_v is the V-band geometric albedo, d is the diameter of an asteroid in km and H_v is the absolute magnitude of the V-band. Albedos of the first three asteroids with minimum spectral deviation from the object are presented in Table 2. As mentioned above, 2000 DP₁₀₇ should be classified as type “M” based on the moderate albedo and the spectral feature.

4. Discussion

For those fast-moving NEAs such as 2000 DP₁₀₇, which approach the earth in a comparatively short period, it is difficult to observe them over a long-time period. However, even if there is only one night of photometry observation, assuming the data are in good precision and the rotation period of the asteroid is short, it is still possible to obtain a credible rotation period at a certain extent, as can be seen from the comparison of our photometric results with those of Pravec et al. (2000).

Although the taxonomic classification of 2000 DP₁₀₇ as an M-type asteroid based on spectrum and albedo is certain, its density of 1.7 g/cm³ from the radar results is at odds with the conventional interpretation that “M” type is dominated by metallic materials and thus has high density.

In this paper, we present a method to derive the spectral type of an asteroid based on a current spectral classification system. The result indicates that asteroid classification based on broad band photometry might not be precise enough, especially for distinguishing between C- and X-type asteroids.

Acknowledgements

The authors thank Prof. Wenzhang Ma, Prof. Wenzhong Liu and Miss Min Guan for their helpful discussions and Xiaoming Teng, Shulin Kong and Jinhu Wang for their assistance with the observations. We also Thank Dr. Alan W. Harris and Dr. Xu Zhou for their valuable comments. This work was supported by the National Natural Science Foundation of China (grant no. 10073012 and no. 19833020). Data reductions of the work are performed in the CAS-PKU Jointly Beijing Astrophysics Center. The 2.16-m

telescope and 0.6-m/0.9-m Schmidt telescope are operated with support from the Chinese Academy of Sciences.

References

- Binzel, R.P., Bus, S.J., Burbine, T.H., Rivkin, A.S., 2001. Announcing the availability of the MIT SMASS and SMASSIR data sets. LPI 32, No. 1633.
- Blythe, M., Shelly, F., Bezpalko, M., Elowitz, M., Huber, R., 2000. 2000 DP₁₀₇, M.P.E.C., 2000-E20.
- Bowell, E., Hapke, B., Domingue, D., Lumme, K., Peltoniemi, J., Harris, A.W., 1989. Application of photometric models to asteroids. In: Asteroids II. University of Arizona Press, Tucson, pp. 524–556.
- Bus, S.J., 1999. Compositional structure in the asteroid belt: results of a spectroscopic survey. Ph.D. Thesis, Massachusetts Institute of Technology.
- Bus, S.J., Binzel, R.P., 2002. Phase II of the small main-belt asteroid spectroscopic survey a feature-based taxonomy. Icarus 158, 146–177.
- Cochran, A.L., Barnes, T.G. III, 1981. Spectrophotometry with a self-scanned silicon photodiode array. I—Instrumentation and reductions. Astrophys. J. Suppl. 45, 73–81.
- Fan, X., Burstein, D., Chen, J.S., Zhu, J., Jiang, Z., Wu, H., Yan, H., Zheng, Z., Zhou, X., Fang, L.Z., Chen, F., Deng, Z., Chu, Y., Hester, J.J., Windhorst, A., Li, Y., Lu, P., Sun, W.H., Chen, W.P., Tsay, W.S., Chiueh, T.H., Chou, C.K., Ko, C.M., Lin, T.C., Guo, H.J., Byun, Y.I., 1996. Deep wide-field spectrophotometry of the open cluster M67. AJ 112, 628.
- Hardorp, J., 1978. The sun among the stars. I—A search for solar spectral analogs. Acta. Astron. 63, 383–390.
- Harris, A.W., Lupishko, D.F., 1989. Photometric lightcurve observations and reduction techniques. In: Asteroids II. University of Arizona Press, Tucson, pp. 39–53.
- Luu, J.X., Jewitt, D.C., 1999. Charge-coupled device spectra of asteroids. I—Near-earth and 3:1 resonance asteroids. AJ 99, 1985–2011.
- Margot, J.L., Nolan, M.C., Benner, L.A.M., Ostro, S.J., Jurgens, R.F., Slade, M.A., Giorgini, J.D., Campbell, D.B., 2000. Satellite of minor planet. IAU Circular No. 7503.
- Margot, J.L., Nolan, M.C., Benner, L.A.M., Ostro, S.J., Jurgens, R.F., Giorgini, J.D., Slade, M.A., Campbell, D.B., 2002. Binary asteroids in the near-earth object population. Science 296, 1445–1448.
- Ostro, S.J., Margot, J.L., Nolan, M.C., Benner, L.A.M., Jurgens, R.F., Giorgini, J.D., 2000. 2000 DP₁₀₇ IAU Circular, No. 7496.
- Pravec, P., Kusnirakv, P., Hicks, M., Holliday, B., Warner, B., 2000. 2000 DP₁₀₇ IAU Circular, No. 7504.
- Richmond, M.W., Droegge, T.F., Gombert, G., Gutzwiller, M., Henden, A.A., Albertson, C., Beser, N., Molhant, N., Johnson, H., 2000. TASS Mark III photometric survey of the celestial equator. PASP 112, 397–408.
- Silva, D.R., Cornell, M., 1992. A new library of stellar optical spectra. Astrophys. J. Suppl. 81, 865–881.
- Stellingwerf, R.F., 1978. Period determination using phase dispersion minimization. Astrophys. J. 224, 953–960.
- Tholen, D.J., 1984. Asteroid taxonomy from cluster analysis of photometry. Ph.D. Thesis, Arizona University, Tucson.
- Tholen, D.J., 1989. Asteroid taxonomic classifications. In: Asteroids II. University of Arizona Press, Tucson, pp. 1139–1150.
- Xu, S., 1994. CCD Photometry and spectroscopy of small main-asteroids. Ph.D. Thesis., Massachusetts Institute of Technology.
- Xu, S., Binzel, R.P., Burbine, T.H., Bus, S.J., 1995. Small main-belt asteroid spectroscopic survey: initial results. Icarus 115, 1–35.
- Zheng, Z.Y., Shang, Z.H., Su, H.J., Burstein, D., Chen, J.S., Deng, Z.G., Byun, Y.I., Chen, R., Chen, W.P., Deng, L.C., Fan, X.H., Fang, L.Z., Hester, J.J., Jiang, Z.J., Li, Y., Lin, W.P., Sun, W.H., Tsay, W.S., Windhorst, R.A., Wu, H., Xia, X.Y., Xu, W., Xue, S.J., Yan, H.J., Zheng, Z., Zhou, X., Zhu, J., Zou, Z.L., Lu, P., 1999. Deep intermediate-band surface photometry of NGC 5907. AJ 117, 2757–2780.